




Fluid Therapy For Wildlife

Rebekah Weiss, CET, CWR
Aves Wildlife Alliance
Neenah, Wisconsin



Outline


- The Importance of Fluids
- What is Dehydration?
 - Assessing Dehydration
 - Clinical Signs of Dehydration
- Fluid Therapy
 - Calculating Fluid Requirements
 - Fluid Types
 - Routes of Administration
- Summary



The Importance of Fluids


“We never know the worth of water till the well is dry.”

– Thomas Fuller



Why is water important?


- **Living things are made of water.**
 - 60% of an adult’s body weight is water.
 - 80-90 % in young animals.
- Water in the form of blood is the transport mechanism for the bodies required nutrients and also for the removal of waste.
- Without adequate water hormones, medications, and chemical messages can not reach their destinations.



Water Movement in the Body

Definitions


- **Solute:** is a substance dissolved in another substance
- **Solvent:** is a substance that dissolves a solute (a chemically different liquid, solid or gas), resulting in a solution
- **Osmosis:** the net movement of solvent molecules through a partially permeable membrane into a region of higher solute concentration, in order to equalize the solute concentrations on the two sides



Hydration: A balance of gains and losses


To maintain adequate hydration the body is constantly moving water into or out of one of the three major reservoirs.

- Intravascular fluid: in the vessels (5-7%)
- Interstitial fluid: baths non-blood cells (25%)
- Intracellular fluid: inside the cell (60%)



What's the normal rate of fluid loss?

Occurs roughly at a daily rate of 60ml/kg of body weight or 6% of the animal's weight each day.



Sources of Water	Losses of Water
Ingestion of water	Elimination (urine/feces)
Water in food	Respiration
Water produced in metabolism	Evaporation

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

7

What is Dehydration?

Dehydration is the loss of water and salts essential for normal body function.

AVES Wildlife Alliance

Dehydration: The Common Causes

<p>Decreased Intake</p> <ul style="list-style-type: none"> • Systemic illness • Starvation • Injury 	<p>Increased Loss</p> <ul style="list-style-type: none"> • Bleeding (trauma) • Polyuria (urine) • Diarrhea • Vomiting • Panting • Burns
---	--

AVES Wildlife Alliance


Fluid Therapy for Wildlife - 2/28/2017

9

Dehydration: The Results

What happens to you when you are dehydrated?

- Thirst/dry mouth
- Loss of appetite
- Dry skin
- Constipation
- Decreased urination
- Weakness
- Headache
- Muscle cramps
- Dizziness
- Decreased blood pressure



AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

10

Assessing Dehydration

How to tell when your patient is more than thirsty.

AVES Wildlife Alliance


Assessing Dehydration: Skin Tenting

- **Definition:** The time it takes for skin pulled away from the body to fall back into a normal resting position.
- Where to Pinch
 - **Mammals:** skin between shoulder blades.
 - **Birds:** skin over the toes or eyelids.
 - **Reptiles:** skin tenting is not a reliable test.
- Dehydration is present when it takes longer than **one second** for the skin to return to a normal resting position.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

12



Assessing dehydration by the skin tent method on a Virginia Opossum

AVES Wildlife Alliance

Fluid Therapy for Wildlife 2/28/2017

13



Dehydrated American Robin hatchling. Note the wrinkled abdominal skin.

AVES Wildlife Alliance

Fluid Therapy for Wildlife 2/28/2017

14


Assessing Dehydration: Capillary Refill Time

- **Definition:** The time it takes for the capillaries to refill after applying enough pressure to blanch the surrounding area.
- Where to Apply Blanching Pressure
 - **Mammals:** perform on the gums (can be difficult with aggressive patients)
 - **Birds:** use the basilic/ulnar vein (inside of elbow)
 - **Reptiles:** difficult
- A Capillary Refill Time (CRT) of greater than 2 seconds is indicative of dehydration greater than 7%.

AVES Wildlife Alliance

Fluid Therapy for Wildlife 2/28/2017

15



Assessing dehydration by the CRT method on a Virginia Opossum.

AVES Wildlife Alliance

Fluid Therapy for Wildlife 2/28/2017

16



Ulnar vein on a Great Horned Owl. CRT assessment site.

AVES Wildlife Alliance

Fluid Therapy for Wildlife 2/28/2017

17


Assessing Dehydration: Mucous Membranes

- **Definition:** Visually examining the color and texture of the mucous membranes.
- Where to look
 - Can evaluate oral membranes, eye conjunctiva, vulva or cloaca.
- What is normal
 - Pale pink to pink membrane color, but know your species not all animals have pink membranes.
 - Glistening and moist.
 - If membranes are pale and tacky or the animal has thick mucous strands in mouth they are dehydrated.

AVES Wildlife Alliance

Fluid Therapy for Wildlife 2/28/2017

18



Clinical Signs of Dehydration

In most cases with wild animals, particularly if there is trauma or disease, it is safe to assume there is at least 5% dehydration.

Clinical Signs of Dehydration: Birds

Clinical Signs	Percent Dehydration
No obvious signs, but assume all injured or orphaned birds have a fluid deficit.	< 5%
Skin may appear tight, especially over the keel. It may also appear dry and wrinkly. The skin forms a temporary tent if pulled up. Eyes look dull and eyelids tent. Inside of mouth is tacky, mucous strands.	5-10%
Inside of mouth is dry. Extremities are cold. Skin remains tented when pulled up. Heartbeat is rapid. Bird looks ill, listless and depressed.	10-15%

Clinical Signs of Dehydration: Mammals

Clinical Signs	Percent Dehydration
No obvious signs, but assume all injured or orphaned mammals have a fluid deficit.	< 5%
Skin appears tight; forms temporary tent if pulled up. Inside of mouth is dry. Decreased capillary refill time. Eyes starting to appear dry and sunken. Urine is concentrated and decreased volume.	5-7%
Pulse is weak and rapid. Eyes are sunken. Extremities are cold. Skin remains tented when pulled up. Mucous membranes are pale. Life threatening.	7-10%

Clinical Signs of Dehydration: Reptiles

Clinical Signs	Percent Dehydration
Increased thirst, slight lethargy, decreased urates.	3%
Increased thirst, anorexia and dullness. Dry loose wrinkled skin; slowly returns to normal when tented. Dull eyes. Dry, often sticky mucous oral membranes. Will start to see dysecdysis (abnormal shedding.)	7%
Dull to comatose. Skin remains tented after pinching. Animal feels light. Dry mucous membranes. Sunken eyeballs. No urates.	10%


Laboratory Evaluation

- Hematocrit (PCV)
- Total plasma protein concentration
- Urine specific gravity (mammals)
- All 3 can be elevated with dehydration.

Test	Birds	Mammals	Reptiles
PCV	35-55%	35-55%	20-40%
Total Protein	3.5-5.5 g/dl	5.5-7.5 g/dl	3.0-8.0 g/dl


Fluid Therapy

The regulation of water balance in a patient by careful measurement of fluid intake against daily losses.



Cautionary Note

- Fluid Therapy is an inexact science.
- The fluid in the body is in a continuous state of flux through normal physiologic processes.
- There are many diverging opinions regarding the type of fluid, volumes and rates of fluid administration and when and how to determine when enough fluid has been administered.
- **The point is that fluid therapy must be reassessed at frequent intervals and adjusted to obtain the maximum benefit and results.**





Goals of Fluid Therapy

- **Maintenance Fluids**
 - Provide daily needs
- **Fluid Deficite**
 - Correct existing deficits
- **Ongoing Losses**
 - To replace ongoing losses from bleeding, diarrhea, kidney problems, etc.




Additional Benefits of Fluid Therapy

- Strengthens patient
- Faster recovery
- Warms patient
- Increases circulation
- Detoxifies –increases blood passing through the liver and kidneys
- Stimulates the gastrointestinal tract


Getting Started

- Once the degree of dehydration is determined the next step is to develop a plan that includes:
 - **Total fluid requirement**
 - **Fluid type**
 - **Administration route**





Calculating Fluid Requirement

Math can be your friend.
It is time to learn how.



Calculating Fluid Requirements



Fluid Therapy for Wildlife - 2/28/2017

31



Pieces of the Fluid Calculation Puzzle

Common Conversions

- 1 kg = 1000 g
- 1 lb = 454 g
- 1 ml = 1 cc

Maintenance Fluids

- 6% of body weight

Fluid Deficit

- % dehydration

Ongoing Losses

- Estimate of ongoing losses from bleeding, diarrhea, kidney problems, etc.

Fluid Therapy for Wildlife - 2/28/2017

32



Total Fluid Requirement: Step One

Patient is a 1.2 kg Red-tailed Hawk that is 8% dehydrated and is losing 7 ml of blood per day as a result of an actively bleeding wound

Step one: Convert the weight to g
1 kg = 1000 g

$$1.2 \text{ kg} \times 1000 \text{ g} = 1200 \text{ g}$$

Fluid Therapy for Wildlife - 2/28/2017

33



Total Fluid Requirement: Step 2

Step Two: Multiply the body weight in g by the percentage of required daily maintenance fluids plus the fluid deficit

$$\text{BW (g)} \times (\text{Maintenance Fluids} + \text{Fluid Deficit}) =$$

$$\text{BW (g)} \times (\text{Maintenance (6\%)} + \text{Fluid Deficit (8\%)}) =$$

$$1200 \text{ g} \times (6\% + 8\%) =$$

$$14\% = 0.14$$

$$1200 \text{ g} \times (0.14) = 168 \text{ ml}$$

Fluid Therapy for Wildlife - 2/28/2017

34



Total Fluid Requirement: Step 3

Step Three: Add the ongoing losses to the answer from step two

7 ml of daily blood loss

Answer from step two + ongoing losses =

$$168 \text{ ml} + 7 \text{ ml} = 175 \text{ ml}$$

175 ml or cc of fluids per 24 hours

Fluid Therapy for Wildlife - 2/28/2017

35



Total Fluid Requirement: Step 4

Step Four: Calculate how much fluid to administer.

- Normal Crop or Stomach Capacity

$$\text{Body Weight (g)} \times 5\% = \text{Maximum crop/stomach capacity}$$

- Debilitated Crop or Stomach Capacity

$$\text{Body Weight (g)} \times 2.5\% = \text{Debilitated capacity}$$

- Total Subcutaneous (SC) Capacity

$$\text{Body Weight (g)} \times 3\% = \text{Maximum SC capacity}$$

Fluid Therapy for Wildlife - 2/28/2017

36



Total Fluid Requirement: Step 4

Calculate Normal Crop/Stomach Capacity
 Body Weight (g) x 5% = Dehydrated crop/stomach capacity

$$1200\text{g} \times 0.05 (5\%) = 60 \text{ cc or ml}$$

Gavage 60 ml per feeding



37

Total Fluid Requirement: Step 5

Step 5: Determine fluid frequency.

Divide total fluid requirement (175 ml)
 by
 the crop capacity (60 ml)

$$175 \text{ ml} / 60 \text{ ml} = 2.9$$

Gavage 60 ml 3 times daily



38

Creating a Multiday Plan

- Day One
 - Maintenance Fluids
 - Fluid Deficit
 - Ongoing Losses
- Day Two & Beyond
 - Maintenance Fluids
 - Ongoing Losses

Remember that this calculation does not replace ongoing patient assessments and monitoring. Stress of frequent handling and the animal's overall condition must always guide our care.



39

Therapeutic Fluid Types

Do you want that shaken or stirred?




Fluid Types

Definitions

- **Osmotic pressure:** the minimum pressure needed to nullify osmosis.
- **Tonicity:** measure of osmotic pressure, comparison of solutes in two solutions
- **Isotonic:** concentrations of solutes that are comparable to normal body fluids
- **Hypertonic:** higher concentrations of solutes than in normal body fluids
- **Hypotonic:** less concentration of solutes than in normal body fluids




41




42


Hypertonic



Isotonic



Hypotonic




Osmotic pressure on human red blood cells.

20 μm
G. Storz, Comment

Fluid Therapy for Wildlife - 2/28/2017

43



Therapeutic Fluid Types

Oral Rehydration Solutions

- Commercial products for human pediatric or calf diarrheal treatments
- Can be given PO only.

Crystalloids


- Can enter cells and all body fluid compartments.
- Used for replacement or maintenance of blood volume.
- Can be given PO, SQ, IV.

Colloids

- Large molecular weight substances that are restricted to bloodstream.
- More of this fluid remains in the bloodstream.
- Ideal for animals in shock.
- Must be given IV or IO.
 - Hetastarch
 - Oxyglobin
 - Whole blood or plasma

Fluid Therapy for Wildlife - 2/28/2017

44




Fluid Type: Hypertonic

Oral rehydration solutions should be hypertonic and should contain glucose and sodium.


- Pedialyte
- Equalyte
- Dioralyte ORS

Caution
For oral use only!



Fluid Therapy for Wildlife - 2/28/2017

45




Fluid Type: Isotonic

Administer SC or IV. If given PO should be diluted and with glucose added.


- Lactated Ringers Solution
- 0.9% Sodium Chloride
- Normosol-R
- 2.5% Dextrose
- Plasma-LTE

Caution
5% Dextrose is Hypertonic
Do not use SC!




Fluid Therapy for Wildlife - 2/28/2017

46




Fluid Management

- Store fluids in cool, dark place.
- Place date opened on bag/bottle.
- Always check to make sure there is no discoloration or particulate matter floating in fluids prior to use.



Fluid Therapy for Wildlife - 2/28/2017


47



Routes of Administration

The route of administration should be determined by the animal's condition, the volume of fluid needed to be given and the expertise of the administrator.

Fluid Therapy for Wildlife - 2/28/2017



Choosing the Right Route

Route	Advantages	Disadvantages
Oral (PO)	Cheap, very effective, most natural. Helps flush the GI system.	Must be conscious, holding head up and swallowing. Risk of aspiration.
Subcutaneous (SC)	Simple, cheap, effective, can give large bolus to minimize handling.	Minimal absorption if animal is in shock. Must be administered using sterile techniques.
Intravenous (IV) Intraosseous (IO)	Delivers fluid directly into blood volume.	Risk of infection. Stress from catheter placement; can fall out. Risk of fluid overload.
Peritoneal (IP)	Can give large bolus, absorbed slowly over time.	Risk of infection, puncturing an internal organ. Must use correct fluid temperature.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

49

Which Route of Administration is Best?

- In patients with a functional gastrointestinal tract (normal bowel movements), PO fluids are best.
- SQ fluids can be used for dehydrated patients that have diminished GIT function (diarrhea, vomiting) or in patients unable to maintain a normal upright posture.
- When replacing large fluid deficits (critical patients), IV or IO routes are best.
- Fluids should never be administered into body cavities of birds (can enter air sacs and drown the patient).

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

50

Oral Fluids: Gavage


If the gut works, use it.

AVES Wildlife Alliance

Gavage: Step by Step Instructions

Getting Ready: Gavage


- Calculate the gavage volume.
- Measure out the solution in an appropriately sized syringe.
- Select the appropriately sized gavage tube.
 - Should be a larger diameter than the glottis or trachea.



AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

52



Glottis on a Great Horned Owl.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

53




Warm the gavage solution to the approximate temperature of the animal.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017


54



Externally measure the length of tube needed to reach the stomach or crop.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017


55



Mark the tube length with a small piece of tape.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017


56



Lubricate the end of the tube with a water based lubricant.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017


57



Charge the tube with the gavage solution to prevent injecting air.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017


58



Extend the animals head upward and in a bird identify the glottis.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017


59



Insert tube past the glottis to the premeasured tape mark.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017

60




Fluid Therapy for Wildlife - 2/28/2017

AVES Wildlife Alliance

Slowly dispense the gavage solution. Watch the back of the mouth for any fluid.

61



Fluid Therapy for Wildlife - 2/28/2017

AVES Wildlife Alliance

Pinch the tube & withdraw. Slowly let the head come back to a normal position.

62

Subcutaneous Fluids

This simple procedure, if done right, is one of the most critical in a wildlife rehabilitators life saving tool box.

AVES Wildlife Alliance

Sites for Subcutaneous Fluids

Birds

- Inguinal skin fold** (the juncture of the leg and body).
- Interscapular region** (featherless tract on either side of the spine over the shoulder area).
- Never inject into the patagium in the armpit.
- Avoid if patient has subcutaneous emphysema – air bubbles under the skin.

Mammals

- Loose skin along the back of the neck between the shoulder blades.
- Almost any area of loose skin along the back and flanks.

Reptiles

- Inguinal or axillary area (turtles).
- On either side of the spine in the mid-third of the body (snakes).

Fluid Therapy for Wildlife - 2/28/2017


AVES Wildlife Alliance

64

SC Fluids: Step by Step Instructions

Getting Ready: SC Fluids

- Calculate the fluid volume.
- Using sterile technique, draw out the fluids into an appropriately sized sterile syringe.
- Select the appropriately sized sterile needle.




Fluid Therapy for Wildlife - 2/28/2017

AVES Wildlife Alliance

65

Fluid Management

- Fluids should be warmed to roughly body temperature prior to administration.



Species	Fluid Temperature
Birds	100 –104° F
Mammals	96–100° F
Marsupials	94 –98° F
Reptiles	86 –89.5° F

Fluid Therapy for Wildlife - 2/28/2017

AVES Wildlife Alliance

66




Locate injection site. Vest on a mammal.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

67




Locate injection site. Inguinal area on a bird.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

68




Wet the fur or feathers with alcohol to visualize the injection site.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

69




Make sure the needle is bevel side up.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

70




Angle the syringe so that it is parallel to the skin.

AVES Wildlife Alliance

Fluid Therapy for Wildlife - 2/28/2017

71




Insert needle under the skin, pull back on the plunger to check for a blood flash.

AVES Wildlife Alliance


Fluid Therapy for Wildlife - 2/28/2017

72



A fluid blister will form as the fluids are slowly expelled.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017
73




If the skin becomes very taut, move to a new site to administer remaining fluid.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017
74

Signs of Rehydration


- Increased weight
- Increased urination
- Eyes brighter
- Gums less tacky
- Improved attitude
- Increased appetite
- Improved skin elasticity
- Stronger pulse
 - Increase blood pressure
 - Normal CRT
- Decreased PCV/TP



AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017
75

When to Stop

- Once an animal is eating and drinking readily on its own.
- Hydration has been restored and the patient is able to maintain hydration without support.
- May want to taper fluid therapy by decreasing the amount administered by 25% per day.



AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017
76

Summary

- Maintaining an adequate fluid balance in the body is critical for normal body function.
- As rehabilitators we are responsible for helping maintain that balance for our wildlife patients.
- You can save lives by understanding how to calculate fluid volumes, choose the correct fluid type and administer fluids.

AVES Wildlife Alliance
Fluid Therapy for Wildlife - 2/28/2017
77

Credits

Rebekah Weiss, CET, CWR graduated from the University of Wisconsin Stevens Point with a B.S. in Wildlife Ecology, Animal Physiology, and Captive Wildlife Management. She is a Wisconsin State and USFWS Licensed Wildlife Rehabilitator and Educator.

.....

Founded in 2009 Aves Wildlife Alliance is a 501c3 non-profit organization committed to conserving and restoring native wildlife through innovation and excellence in community outreach and wildlife rehabilitation.

Aves Wildlife Alliance
351 Muttart Road, Neenah, WI 54956
920.716.2790
rweiss@aveswildlife.org
www.aveswildlife.org

AVES Wildlife Alliance